

2015 AMC 12A Problems

1. What is the value of $(2^0 - 1 + 5^2 + 0)^{-1} \times 5$?

- (A) -125 (B) -120 (C) $\frac{1}{5}$ (D) $\frac{5}{24}$ (E) 25

2. Two of the three sides of a triangle are 20 and 15. Which of the following numbers is not a possible perimeter of the triangle?

- (A) 52 (B) 57 (C) 62 (D) 67 (E) 72

3. Mr. Patrick teaches math to 15 students. He was grading tests and found that when he graded everyone's test except Payton's, the average grade for the class was 80. After he graded Payton's test, the class average became 81. What was Payton's score on the test?

- (A) 81 (B) 85 (C) 91 (D) 94 (E) 95

4. The sum of two positive numbers is 5 times their difference. What is the ratio of the larger number to the smaller?

- (A) $\frac{5}{4}$ (B) $\frac{3}{2}$ (C) $\frac{9}{5}$ (D) 2 (E) $\frac{5}{2}$

5. Amelia needs to estimate the quantity $\frac{a}{b} - c$, where a , b , and c are large positive integers. She rounds each of the integers so that the calculation will be easier to do mentally. In which of these situations will her answer necessarily be greater than the exact value of $\frac{a}{b} - c$?

- (A) She rounds all three numbers up.
(B) She rounds a and b up, and she rounds c down.
(C) She rounds a and c up, and she rounds b down.
(D) She rounds a up, and she rounds b and c down.
(E) She rounds c up, and she rounds a and b down.

6. Two years ago Pete was three times as old as his cousin Claire. Two years before that, Pete was four times as old as Claire. In how many years will the ratio of their ages be 2 : 1?

- (A) 2 (B) 4 (C) 5 (D) 6 (E) 8

7. Two right circular cylinders have the same volume. The radius of the second cylinder is 10% more than the radius of the first. What is the relationship between the heights of the two cylinders?
- (A) The second height is 10% less than the first.
(B) The first height is 10% more than the second.
(C) The second height is 21% less than the first.
(D) The first height is 21% more than the second.
(E) The second height is 80% of the first.
8. The ratio of the length to the width of a rectangle is 4 : 3. If the rectangle has diagonal of length d , then the area may be expressed as kd^2 for some constant k . What is k ?
- (A) $\frac{2}{7}$ (B) $\frac{3}{7}$ (C) $\frac{12}{25}$ (D) $\frac{16}{25}$ (E) $\frac{3}{4}$
9. A box contains 2 red marbles, 2 green marbles, and 2 yellow marbles. Carol takes 2 marbles from the box at random; then Claudia takes 2 of the remaining marbles at random; and then Cheryl takes the last two marbles. What is the probability that Cheryl gets 2 marbles of the same color?
- (A) $\frac{1}{10}$ (B) $\frac{1}{6}$ (C) $\frac{1}{5}$ (D) $\frac{1}{3}$ (E) $\frac{1}{2}$
10. Integers x and y with $x > y > 0$ satisfy $x + y + xy = 80$. What is x ?
- (A) 8 (B) 10 (C) 15 (D) 18 (E) 26
11. On a sheet of paper, Isabella draws a circle of radius 2, a circle of radius 3, and all possible lines simultaneously tangent to both circles. Isabella notices that she has drawn exactly $k \geq 0$ lines. How many different values of k are possible?
- (A) 2 (B) 3 (C) 4 (D) 5 (E) 6
12. The parabolas $y = ax^2 - 2$ and $y = 4 - bx^2$ intersect the coordinate axes in exactly four points, and these four points are the vertices of a kite of area 12. What is $a + b$?
- (A) 1 (B) 1.5 (C) 2 (D) 2.5 (E) 3
13. A league with 12 teams holds a round-robin tournament, with each team playing every other team once. Games either end with one team victorious or else end in a draw. A team scores 2 points for every game it wins and 1 point for every game it draws. Which of the following is **not** a true statement about the list of 12 scores?
- (A) There must be an even number of odd scores.
(B) There must be an even number of even scores.

(C) There cannot be two scores of 0.

(D) The sum of the scores must be at least 100.

(E) The highest score must be at least 12.

14. What is the value of a for which $\frac{1}{\log_2 a} + \frac{1}{\log_3 a} + \frac{1}{\log_4 a} = 1$?

(A) 9 (B) 12 (C) 18 (D) 24 (E) 36

15. What is the minimum number of digits to the right of the decimal point needed to express the fraction $\frac{123456789}{2^{26} \cdot 5^4}$ as a decimal?

(A) 4 (B) 22 (C) 26 (D) 30 (E) 104

16. Tetrahedron $ABCD$ has $AB = 5$, $AC = 3$, $BC = 4$, $BD = 4$, $AD = 3$, and $CD = \frac{12}{5}\sqrt{2}$. What is the volume of the tetrahedron?

(A) $3\sqrt{2}$ (B) $2\sqrt{5}$ (C) $\frac{24}{5}$ (D) $3\sqrt{3}$ (E) $\frac{24}{5}\sqrt{2}$

17. Eight people are sitting around a circular table, each holding a fair coin. All eight people flip their coins and those who flip heads stand while those who flip tails remain seated. What is the probability that no two adjacent people will stand?

(A) $\frac{47}{256}$ (B) $\frac{3}{16}$ (C) $\frac{49}{256}$ (D) $\frac{25}{128}$ (E) $\frac{51}{256}$

18. The zeroes of the function $f(x) = x^2 - ax + 2a$ are integers. What is the sum of all possible values of a ?

(A) 7 (B) 8 (C) 16 (D) 17 (E) 18

19. For some positive integers p , there is a quadrilateral $ABCD$ with positive integer side lengths, perimeter p , right angles at B and C , $AB = 2$, and $CD = AD$. How many different values of $p < 2015$ are possible?

(A) 30 (B) 31 (C) 61 (D) 62 (E) 63

20. Isosceles triangles T and T' are not congruent but have the same area and the same perimeter. The sides of T have lengths 5, 5, and 8, while those of T' have lengths a , a , and b . Which of the following numbers is closest to b ?

(A) 3 (B) 4 (C) 5 (D) 6 (E) 8

21. A circle of radius r passes through both foci of, and exactly four points on, the ellipse with equation $x^2 + 16y^2 = 16$. The set of all possible values of r is an interval $[a, b)$. What is $a + b$?

(A) $5\sqrt{2} + 4$ (B) $\sqrt{17} + 7$ (C) $6\sqrt{2} + 3$ (D) $\sqrt{15} + 8$ (E) 12

22. For each positive integer n , let $S(n)$ be the number of sequences of length n consisting solely of the letters A and B , with no more than three A s in a row and no more than three B s in a row. What is the remainder when $S(2015)$ is divided by 12?

- (A) 0 (B) 4 (C) 6 (D) 8 (E) 10

23. Let S be a square of side length 1. Two points are chosen independently at random on the sides of S . The probability that the straight-line distance between the points is at least $\frac{1}{2}$ is $\frac{a-b\pi}{c}$, where a , b , and c are positive integers and $\gcd(a, b, c) = 1$. What is $a + b + c$?

- (A) 59 (B) 60 (C) 61 (D) 62 (E) 63

24. Rational numbers a and b are chosen at random among all rational numbers in the interval $[0, 2)$ that can be written as fractions $\frac{n}{d}$ where n and d are integers with $1 \leq d \leq 5$. What is the probability that


$$(\cos(a\pi) + i \sin(b\pi))^4$$

is a real number?

- (A) $\frac{3}{50}$ (B) $\frac{4}{25}$ (C) $\frac{41}{200}$ (D) $\frac{6}{25}$ (E) $\frac{13}{50}$

25. A collection of circles in the upper half-plane, all tangent to the x -axis, is constructed in layers as follows. Layer L_0 consists of two circles of radii 70^2 and 73^2 that are externally tangent. For $k \geq 1$, the circles in $\bigcup_{j=0}^{k-1} L_j$ are ordered according to their points of tangency with the x -axis. For every pair of consecutive circles in this order, a new circle is constructed externally tangent to each of the two circles in the pair. Layer L_k consists of the 2^{k-1} circles constructed in this way. Let $S = \bigcup_{j=0}^6 L_j$, and for every circle C denote by $r(C)$ its radius. What is

$$\sum_{C \in S} \frac{1}{\sqrt{r(C)}}?$$



- (A) $\frac{286}{35}$ (B) $\frac{583}{70}$ (C) $\frac{715}{73}$ (D) $\frac{143}{14}$ (E) $\frac{1573}{146}$